
SureGuard[®] SC



Residual Control that Lasts – Now in a Convenient Liquid Formula

New from Nufarm, liquid SureGuard® SC provides an exceptional value that allows you to offer more and make more - a perfect solution for your customers. SureGuard SC delivers the same long lasting residual control of grass and broadleaf weeds that you've come to expect from SureGuard in a new convenient liquid formulation. Also, SureGuard SC enhances the speed and spectrum of glyphosate when used as a tank mix partner, even on glyphosate resistant weeds.

TOUGH CONTROL, QUICK

- LABEL UPDATE: Now approved for aquatic and bareground use sites
- Longest residual pre-emergent control
- Convenient easy to mix and apply patent pending liquid formulation
- Broad spectrum including tough-to-control weeds
- Fast and effective knockdown when tank mixed with glyphosate, even on glyphosate-resistant weeds
- Non-staining, spray formulation
- No yellow color, odor, or concerns with off-target movement and volatility
- One product for weed control in landscapes, bareground, cracks and crevices, and dormant Bermudagrass

GREATER EASE, SAME POWER

USE RATES ARE EQUIVALENT TO ORIGINAL DRY FORMULATION:
1 liquid fl oz SureGuard SC = 1 dry oz SureGuard WDG

See label for complete directions for use

EPA REG. NO.	71368-114
ACTIVE INGR.	flumioxazin (44%)
FORMULATION	liquid SC
CHEM. FAMILY	N-phenylphthalimide
HRAC NO.	14
SIGNAL WORD	no signal word required
RES. USE	no
PACKAGE SIZE	8 x 1 pt, 4 x 64 fl oz case







LANDSCAPE ORNAMENTALS

PRE-EMERGE
POST-EMERGE

AQUATIC



SureGuard SC

KEY USES

Dormant Bermudagrass on residential sites, golf courses, and sod farms

Landscape ornamentals

Non-bearing fruit and nut trees and vines Non-crop areas: around buildings, fencerows, gravel surfaces, driveways, and ground mats Slow moving or quiescent aquatic sites

KEY WEEDS CONTROLLED

(See product label for complete list)

Bittercress

Barnyardgrass*

Bluegrass*

Chickweed

Crabgrass*

Dandelion*

Eclipta

Field pennycress

Florida beggarweed

Florida pusley

Foxtail*

Goosegrass*

Groundsel

Kochia

Lambsquarters

Mallow

Morningglory

Nightshade

Palmer amaranth

Pennsylvania smartweed

Piaweed

Pineapple-weed*

Prickly sida (teaweed)

Puncturevine

Ragweed

Redmaids

Russian thistle

Shepherd's-purse

Spiny amaranth

Spurge

Velvetleaf

Waterhemp

Yellow rocket

*Pre-emerge only

AQUATIC WEEDS

Alligator weed

Duckweed

Frog's bit

Water fern

Water lettuce Watermeal

USE SUMMARY

AQUATIC AREAS

Apply this product as a broadcast spray at 6 to 12 fl oz of formulated product per acre plus an adjuvant approved for use in aquatics.

Apply this product in a minimum of 30 gallons of water per acre to all areas of the water body where weeds exist. If a second application is required to provide control, it is recommended that a treatment be made once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

Application of this product during early morning hours may enhance weed control. When applying to densely packed actively growing surface weeds, ensure adequate coverage.

This product may be tank mixed with 2,4-D, diquat, glyphosate or other registered foliar applied herbicides for enhanced control of floating and emergent weeds.

WOODY LANDSCAPE ORNAMENTALS AND BAREGROUND AREAS

Application of SureGuard® SC in the vicinity of ornamental plants is limited to directed sprays around well established woody shrubs and conifer and deciduous trees listed in the product label. SureGuard SC may also be applied to maintain bareground in non-crop areas as listed in the product label. Do not apply this product within any enclosed structure in residential or commercial properties.

Pre-emergence Weed Control: Mix 0.18 - 0.27 fluid ounces of this product per gallon of spray solution and apply 1 gallon of spray solution to 1,000 square feet prior to weed germination. Apply to weed free soil, mulch or gravel surfaces. Moisture is necessary to activate this product on soil for residual weed control.

Post-emergence Weed Control: Mix 0.18 - 0.27 fluid ounces of this product per gallon and apply 1 gallon of spray solution to 1,000 square feet to actively growing weeds. Tank mixing this product with glyphosate will increase the spectrum of post-emergent weed control and provide faster post-emergent weed control than glyphosate alone.

Established landscape ornamentals have shown tolerance to applications of this product only when applied to the soil at the base of the plant and sprays do not directly contact or drift onto desirable plants. For maximum plant safety when using around desirable ornamentals, direct applications of this product (or this product plus glyphosate) toward the soil and leave a sufficient non-treated buffer to ensure spray solution does not contact desired plants.

DORMANT BERMUDAGRASS (Only for use in certain states, see product label)

SureGuard SC may be applied as a single or split application to well established dormant Bermudagrass. It will provide pre-emergence and early post-emergence control of annual bluegrass, chickweed, henbit and other winter annual weeds listed in the product label and pre-emergence control of crabgrass, goosegrass and other summer annual weeds listed in the product label. Application of this product to actively growing turfgrass or during green-up may cause unacceptable injury.

Broadcast Application: Apply 8 - 12 fluid ounces of this product per broadcast acre as a preemergence application. If weeds are present at the time of application apply this product plus an adjuvant (0.25% v/v non-ionic surfactant). Make post-emergence applications when weeds are actively growing and no larger than 2" in height. Thorough spray coverage is necessary to maximize the post-emergence activity. When applied after weed germination, this product will provide pre-emergence and post-emergence control of broadleaf weeds and grasses listed in the product label. Post-emergence weed control may be more effective on certain weed species, and may not control mature, stressed or hardened off weeds that are not actively growing at the time of application.

See label for complete use directions, restrictions and limitations.

